

Finite-time future singularities in modified gravity

Kazuharu Bamba*

Department of Physics, National Tsing Hua University, Hsinchu, Taiwan 300

We review finite-time future singularities in modified gravity. We reconstruct an explicit model of modified gravity realizing a crossing of the phantom divide and show that the Big Rip singularity appears in the modified gravitational theory. It is also demonstrated that the (finite-time) Big Rip singularity in the modified gravity is transformed into the infinite-time singularity in the corresponding scalar field theory obtained through the conformal transformation. Furthermore, we study several models of modified gravity which produce accelerating cosmologies ending at the finite-time future singularities of all four known types.

I. INTRODUCTION

Recent observations confirmed that the current expansion of the universe is accelerating. There are two approaches to explain the current accelerated expansion of the universe. One is to introduce some unknown matter, which is called “dark energy” in the framework of general relativity. The other is to modify the gravitational theory, e.g., to study the action described by an arbitrary function of the scalar curvature R . This is called “ $F(R)$ gravity”, where $F(R)$ is an arbitrary function of the scalar curvature R (for reviews, see [1, 2]).

According to the recent various observational data, there exists the possibility that the effective equation of state (EoS) parameter, which is the ratio of the effective pressure of the universe to the effective energy density of it, evolves from larger than -1 (non-phantom phase) to less than -1 (phantom one, in which superacceleration is realized), namely, crosses -1 (the phantom divide) currently or in near future. Various attempts to realize the crossing of the phantom divide have been made in the framework of general relativity. Recently, a crossing of the phantom divide in modified gravity has also been investigated [1, 3, 4, 5]. Moreover, it is known that modified gravity may lead to the effective phantom/quintessence phase [1], while the phantom/quintessence-dominated universe may end up with finite-time future singularities, which can be categorized into four types [6].

In the present article, we review finite-time future singularities in modified gravity. Following the considerations in Ref. [5], we reconstruct an explicit model of modified gravity realizing a crossing of the phantom divide by using the reconstruction method proposed in Refs. [7, 8]. We show that the Big Rip singularity appears in this modified gravitational theory, whereas that the (finite-time) Big Rip singularity in the modified gravity is transformed into the infinite-time singularity in the corresponding scalar field theory. Next, following the investigations in Ref. [9], we explore several examples of $F(R)$ gravity which predict the accelerating cosmological solutions ending at the finite-time future singularities. It is demonstrated that not only the Big Rip but other three types of the finite-time future singularities may appear.

This article is organized as follows. In Sec. II we explain the reconstruction method of modified gravity [7, 8]. Using this method, we reconstruct an explicit model of modified gravity in which a crossing of the phantom divide can be realized. We also consider the corresponding scalar field theory. In Sec. III we present several models of $F(R)$ gravity which predict accelerating cosmologies ending at the finite-time future singularities by using the reconstruction method. Finally, summary is given in Sec. IV. We use units in which $k_B = c = \hbar = 1$ and denote the gravitational constant $8\pi G$ by κ^2 , so that $\kappa^2 \equiv 8\pi/M_{\text{Pl}}^2$, where $M_{\text{Pl}} = G^{-1/2} = 1.2 \times 10^{19} \text{GeV}$ is the Planck mass.

II. MODIFIED GRAVITATIONAL THEORY REALIZING A CROSSING OF THE PHANTOM DIVIDE

We investigate modified gravity in which a crossing of the phantom divide can be realized by using the reconstruction method.

A. Reconstruction of modified gravity

First, we review the reconstruction method of modified gravity proposed in Refs. [7, 8] (for the related study of reconstruction in $F(R)$ gravity, see [10]).

* E-mail address: bamba“at”phys.nthu.edu.tw

The action of $F(R)$ gravity with general matter is given by

$$S = \int d^4x \sqrt{-g} \left[\frac{F(R)}{2\kappa^2} + \mathcal{L}_{\text{matter}} \right], \quad (2.1)$$

where g is the determinant of the metric tensor $g_{\mu\nu}$ and $\mathcal{L}_{\text{matter}}$ is the matter Lagrangian.

The action (2.1) can be rewritten to the following form by using proper functions $P(\phi)$ and $Q(\phi)$ of a scalar field ϕ :

$$S = \int d^4x \sqrt{-g} \left\{ \frac{1}{2\kappa^2} [P(\phi)R + Q(\phi)] + \mathcal{L}_{\text{matter}} \right\}. \quad (2.2)$$

The scalar field ϕ may be regarded as an auxiliary scalar field because ϕ has no kinetic term. Taking the variation of the action (2.2) with respect to ϕ , we obtain

$$0 = \frac{dP(\phi)}{d\phi} R + \frac{dQ(\phi)}{d\phi}, \quad (2.3)$$

which may be solved with respect to ϕ as $\phi = \phi(R)$. Substituting $\phi = \phi(R)$ into the action (2.2), we find that the expression of $F(R)$ in the action of $F(R)$ gravity in Eq. (2.1) is given by

$$F(R) = P(\phi(R))R + Q(\phi(R)). \quad (2.4)$$

Taking the variation of the action (2.2) with respect to the metric $g_{\mu\nu}$, we find that the field equation of modified gravity is given by

$$\frac{1}{2}g_{\mu\nu} [P(\phi)R + Q(\phi)] - R_{\mu\nu}P(\phi) - g_{\mu\nu}\square P(\phi) + \nabla_\mu \nabla_\nu P(\phi) + \kappa^2 T_{\mu\nu}^{(\text{matter})} = 0, \quad (2.5)$$

where ∇_μ is the covariant derivative operator associated with $g_{\mu\nu}$, $\square \equiv g^{\mu\nu}\nabla_\mu \nabla_\nu$ is the covariant d'Alembertian for a scalar field, and $T_{\mu\nu}^{(\text{matter})}$ is the contribution to the matter energy-momentum tensor.

We assume the flat Friedmann-Robertson-Walker (FRW) space-time with the metric $ds^2 = -dt^2 + a^2(t)d\mathbf{x}^2$, where $a(t)$ is the scale factor. In this background, the $(\mu, \nu) = (0, 0)$ component and the trace part of the $(\mu, \nu) = (i, j)$ component of Eq. (2.5), where i and j run from 1 to 3, read

$$-6H^2 P(\phi(t)) - Q(\phi(t)) - 6H \frac{dP(\phi(t))}{dt} + 2\kappa^2 \rho = 0, \quad (2.6)$$

$$\text{and} \quad 2 \frac{d^2 P(\phi(t))}{dt^2} + 4H \frac{dP(\phi(t))}{dt} + (4\dot{H} + 6H^2) P(\phi(t)) + Q(\phi(t)) + 2\kappa^2 p = 0, \quad (2.7)$$

respectively, where $H = \dot{a}/a$ is the Hubble parameter. Here, ρ and p are the sum of the energy density and pressure of matters with a constant EoS parameter w_i , respectively, where i denotes some component of the matters.

Eliminating $Q(\phi)$ from Eqs. (2.6) and (2.7), we obtain

$$\frac{d^2 P(\phi(t))}{dt^2} - H \frac{dP(\phi(t))}{dt} + 2\dot{H} P(\phi(t)) + \kappa^2 (\rho + p) = 0. \quad (2.8)$$

We note that the scalar field ϕ may be taken as $\phi = t$ because ϕ can be redefined properly.

We now consider that $a(t)$ is described as $a(t) = \bar{a} \exp(\tilde{g}(t))$, where $\bar{a}$ is a constant and $\tilde{g}(t)$ is a proper function. In this case, Eq. (2.8) is reduced to

$$\frac{d^2 P(\phi)}{d\phi^2} - \frac{d\tilde{g}(\phi)}{d\phi} \frac{dP(\phi)}{d\phi} + 2 \frac{d^2 \tilde{g}(\phi)}{d\phi^2} P(\phi) + \kappa^2 \sum_i (1 + w_i) \bar{\rho}_i \bar{a}^{-3(1+w_i)} \exp[-3(1+w_i)\tilde{g}(\phi)] = 0, \quad (2.9)$$

where $\bar{\rho}_i$ is a constant and we have used $H = d\tilde{g}(\phi)/(d\phi)$. Moreover, it follows from Eq. (2.6) that $Q(\phi)$ is given by

$$Q(\phi) = -6 \left[\frac{d\tilde{g}(\phi)}{d\phi} \right]^2 P(\phi) - 6 \frac{d\tilde{g}(\phi)}{d\phi} \frac{dP(\phi)}{d\phi} + 2\kappa^2 \sum_i \bar{\rho}_i \bar{a}^{-3(1+w_i)} \exp[-3(1+w_i)\tilde{g}(\phi)]. \quad (2.10)$$

Hence, if we obtain the solution of Eq. (2.9) with respect to $P(\phi)$, then we can find $Q(\phi)$. Consequently, using Eq. (2.4), we can reconstruct $F(R)$ gravity for any cosmology expressed by $a(t) = \bar{a} \exp(\tilde{g}(t))$. In Refs. [7, 11, 12],

specific models unifying the sequence: the early-time acceleration, radiation/matter-dominated stage and dark energy epoch have been constructed.

Next, using the above method, we reconstruct an explicit model in which a crossing of the phantom divide can be realized. We start with Eq. (2.9) without matter:

$$\frac{d^2 P(\phi)}{d\phi^2} - \frac{d\tilde{g}(\phi)}{d\phi} \frac{dP(\phi)}{d\phi} + 2 \frac{d^2 \tilde{g}(\phi)}{d\phi^2} P(\phi) = 0. \quad (2.11)$$

By redefining $P(\phi)$ as $P(\phi) = e^{\tilde{g}(\phi)/2} \tilde{p}(\phi)$, Eq. (2.11) is rewritten to

$$\frac{1}{\tilde{p}(\phi)} \frac{d^2 \tilde{p}(\phi)}{d\phi^2} = 25 e^{\tilde{g}(\phi)/10} \frac{d^2 (e^{-\tilde{g}(\phi)/10})}{d\phi^2}. \quad (2.12)$$

We explore the following model: $\tilde{g}(\phi) = -10 \ln \left[(\phi/t_0)^{-\gamma} - C (\phi/t_0)^{\gamma+1} \right]$, where C and γ are positive constants, and t_0 is the present time. In this case, Eq. (2.12) is reduced to $(1/\tilde{p}(\phi)) [d^2 \tilde{p}(\phi)/d\phi^2] = 25\gamma(\gamma+1)/\phi^2$, which can be solved as $\tilde{p}(\phi) = \tilde{p}_+ \phi^{\beta_+} + \tilde{p}_- \phi^{\beta_-}$. Here, $\tilde{p}_\pm$ are arbitrary constants and $\beta_\pm$ are given by $\beta_\pm = [1 \pm \sqrt{1 + 100\gamma(\gamma+1)}]/2$. From the above expression of $\tilde{g}(\phi)$, we find that $\tilde{g}(\phi)$ diverges at finite ϕ when $\phi = t_s \equiv t_0 C^{-1/(2\gamma+1)}$, which tells that there could be the Big Rip singularity at $t = t_s$. We only need to consider the period $0 < t < t_s$ because $\tilde{g}(\phi)$ should be real number. In this case, the Hubble rate $H(t)$ is given by

$$H(t) = \frac{d\tilde{g}(\phi)}{d\phi} = \left(\frac{10}{t_0} \right) \left[\gamma \left(\frac{\phi}{t_0} \right)^{-\gamma-1} + (\gamma+1) C \left(\frac{\phi}{t_0} \right)^\gamma \right] \left[\left(\frac{\phi}{t_0} \right)^{-\gamma} - C \left(\frac{\phi}{t_0} \right)^{\gamma+1} \right]^{-1}, \quad (2.13)$$

where it is taken $\phi = t$.

In the flat FRW background, even for modified gravity described by the action (2.1), the effective energy-density and pressure of the universe are given by $\rho_{\text{eff}} = 3H^2/\kappa^2$ and $p_{\text{eff}} = -(2\dot{H} + 3H^2)/\kappa^2$, respectively. The effective EoS parameter $w_{\text{eff}} = p_{\text{eff}}/\rho_{\text{eff}}$ is defined as $w_{\text{eff}} \equiv -1 - 2\dot{H}/(3H^2)$ [1]. For the case of $H(t)$ in Eq. (2.13), w_{eff} is expressed as $w_{\text{eff}} = -1 + U(t)$, where

$$U(t) \equiv -\frac{2\dot{H}}{3H^2} = -\frac{1}{15} \left[-\gamma + 4\gamma(\gamma+1) \left(\frac{t}{t_s} \right)^{2\gamma+1} + (\gamma+1) \left(\frac{t}{t_s} \right)^{2(2\gamma+1)} \right] \left[\gamma + (\gamma+1) \left(\frac{t}{t_s} \right)^{2\gamma+1} \right]^{-2}. \quad (2.14)$$

For the case of Eq. (2.13), the scalar curvature $R = 6(\dot{H} + 2H^2)$ is expressed as

$$R = 60 \left[\gamma(20\gamma-1) + 44\gamma(\gamma+1) \left(\frac{t}{t_s} \right)^{2\gamma+1} + (\gamma+1)(20\gamma+21) \left(\frac{t}{t_s} \right)^{2(2\gamma+1)} \right] t^{-2} \left[1 - \left(\frac{t}{t_s} \right)^{2\gamma+1} \right]^{-2}. \quad (2.15)$$

In deriving Eqs. (2.14) and (2.15), we have used $t_s = t_0 C^{-1/(2\gamma+1)}$.

When $t \rightarrow 0$, i.e., $t \ll t_s$, $H(t)$ behaves as $H(t) \sim 10\gamma/t$. In this limit, it follows from $w_{\text{eff}} = -1 - 2\dot{H}/(3H^2)$ that the effective EoS parameter is given by $w_{\text{eff}} = -1 + 1/(15\gamma)$. This behavior is identical with that in the Einstein gravity with matter whose EoS parameter is greater than -1 .

On the other hand, when $t \rightarrow t_s$, we find $H(t) \sim 10/(t_s - t)$. In this case, the scale factor is given by $a(t) \sim \bar{a}(t_s - t)^{-10}$. When $t \rightarrow t_s$, therefore, $a \rightarrow \infty$, namely, the Big Rip singularity appears. In this limit, the effective EoS parameter is given by $w_{\text{eff}} = -1 - 1/15 = -16/15$. This behavior is identical with the case in which there is a phantom matter with its EoS parameter being smaller than -1 . Thus, we have obtained an explicit model showing a crossing of the phantom divide.

It follows from $w_{\text{eff}} = -1 - 2\dot{H}/(3H^2)$ that the effective EoS parameter w_{eff} becomes -1 when $\dot{H} = 0$. Solving $w_{\text{eff}} = -1$ with respect to t by using $w_{\text{eff}} = -1 + U(t)$, namely, $U(t) = 0$, we find that the effective EoS parameter crosses the phantom divide at $t = t_c$ given by $t_c = t_s \left[-2\gamma + \sqrt{4\gamma^2 + \gamma/(\gamma+1)} \right]^{1/(2\gamma+1)}$. From Eq. (2.14), we see that when $t < t_c$, $U(t) > 0$ because $\gamma > 0$. Moreover, the time derivative of $U(t)$ is given by

$$\frac{dU(t)}{dt} = -\frac{1}{15} \left[2\gamma(\gamma+1)(2\gamma+1)^2 \right] \left[\gamma + (\gamma+1) \left(\frac{t}{t_s} \right)^{2\gamma+1} \right]^{-3} \left(\frac{1}{t_s} \right) \left(\frac{t}{t_s} \right)^{2\gamma} \left[1 - \left(\frac{t}{t_s} \right)^{2\gamma+1} \right]. \quad (2.16)$$

Eq. (2.16) tells that the relation $dU(t)/(dt) < 0$ is always satisfied because we only consider the period $0 < t < t_s$ as mentioned above. This means that $U(t)$ decreases monotonously. Thus, the value of $U(t)$ evolves from positive to negative. From $w_{\text{eff}} = -1 + U(t)$, we see that the value of w_{eff} crosses -1 . Once the universe enters the phantom phase, it stays in this phase, namely, the value of w_{eff} remains less than -1 , and finally the Big Rip singularity appears because $U(t)$ decreases monotonically.

As a consequence, $P(t)$ is given by $P(t) = \left\{ (t/t_0)^\gamma / \left[1 - (t/t_s)^{2\gamma+1} \right] \right\}^5 \sum_{j=\pm} \tilde{p}_j t^{\beta_j}$. Using Eqs. (2.10), we obtain $Q(t) = -6H \left\{ (t/t_0)^\gamma / \left[1 - (t/t_s)^{2\gamma+1} \right] \right\}^5 \sum_{j=\pm} (3H/2 + \beta_j/t) \tilde{p}_j t^{\beta_j}$.

When $t \rightarrow 0$, from $H(t) \sim 10\gamma/t$, we find $t \sim \sqrt{60\gamma(20\gamma-1)/R}$. In this limit, it follows from Eqs. (2.4) that the form of $F(R)$ is given by

$$F(R) \sim \left\{ \frac{\left[\frac{1}{t_0} \sqrt{60\gamma(20\gamma-1)} R^{-1/2} \right]^\gamma}{1 - \left[\frac{1}{t_s} \sqrt{60\gamma(20\gamma-1)} R^{-1/2} \right]^{2\gamma+1}} \right\}^5 R \sum_{j=\pm} \left\{ \left(\frac{5\gamma-1-\beta_j}{20\gamma-1} \right) \tilde{p}_j [60\gamma(20\gamma-1)]^{\beta_j/2} R^{-\beta_j/2} \right\}. \quad (2.17)$$

On the other hand, when $t \rightarrow t_s$, from $H(t) \sim 10/(t_s - t)$, we obtain $t \sim t_s - 3\sqrt{140/R}$. In this limit, it follows from Eqs. (2.4) that the form of $F(R)$ is given by

$$F(R) \sim \left[\frac{(J/t_0)^\gamma}{1 - (J/t_s)^{2\gamma+1}} \right]^5 R \sum_{j=\pm} \tilde{p}_j J^{\beta_j} \left\{ 1 - \sqrt{\frac{20}{7}} \left[\sqrt{\frac{15}{84}} t_s + (\beta_j - 15) R^{-1/2} \right] \frac{1}{J} \right\}, \quad (2.18)$$

where $J \equiv t_s - 3\sqrt{140/R}$. The above modified gravity may be considered as some approximated form of more realistic, viable theory. From Eq. (2.15), we see that in the above limit the scalar curvature diverges, and that the expression of $F(R)$ in (2.18) can be approximately written as

$$F(R) \approx \frac{2}{7} \left[\frac{1}{3\sqrt{140}(2\gamma+1)} \left(\frac{t_s}{t_0} \right)^\gamma \right]^5 \left(\sum_{j=\pm} \tilde{p}_j t_s^{\beta_j} \right) t_s^5 R^{7/2}. \quad (2.19)$$

B. Corresponding scalar field theory

In this subsection, motivated by the discussion in Ref. [13], we consider the corresponding scalar field theory to modified gravity realizing a crossing of the phantom divide, which is obtained by making the conformal transformation of the modified gravitational theory. (In Ref. [14], the relations between scalar field theories and $F(R)$ gravity have been studied.) By introducing two scalar fields ζ and ξ , we can rewrite the action (2.1) to the following form [1]:

$$S = \int d^4x \sqrt{-g} \left\{ \frac{1}{2\kappa^2} [\xi(R - \zeta) + F(\zeta)] + \mathcal{L}_{\text{matter}} \right\}. \quad (2.20)$$

The form in Eq. (2.20) is reduced to the original expression in Eq. (2.1) by using the equation $\zeta = R$, which is derived by taking variation of the action (2.20) with respect to one auxiliary field ξ . Taking the variation of the form in Eq. (2.20) with respect to the other auxiliary field ζ , we obtain $\xi = F'(\zeta)$, where the prime denotes differentiation with respect to ζ . Substituting this equation into Eq. (2.20) and eliminating ξ from Eq. (2.20), we find

$$S = \int d^4x \sqrt{-g} \left[\frac{1}{2\kappa^2} (F'(\zeta)R + F(\zeta) - F'(\zeta)\zeta) + \mathcal{L}_{\text{matter}} \right]. \quad (2.21)$$

This is the action in the Jordan-frame, in which there exists a non-minimal coupling between $F'(\zeta)$ and the scalar curvature R . We make the following conformal transformation of the action (2.21): $g_{\mu\nu} \rightarrow \hat{g}_{\mu\nu} = e^\sigma g_{\mu\nu}$, where $e^\sigma = F'(\zeta)$. Here, σ is a scalar field and a hat denotes quantities in the Einstein frame, in which the non-minimal coupling between $F'(\zeta)$ and the scalar curvature R in the first term on the right-hand side of Eq. (2.21) disappears. Consequently, the action in the Einstein frame is given by [15, 16]

$$S_E = \int d^4x \sqrt{-\hat{g}} \left[\frac{1}{2\kappa^2} \left(\hat{R} - \frac{3}{2} \hat{g}^{\mu\nu} \partial_\mu \sigma \partial_\nu \sigma - V(\sigma) \right) + e^{-2\sigma} \mathcal{L}_{\text{matter}} \right], \quad (2.22)$$

where $V(\sigma) = e^{-\sigma}\zeta(\sigma) - e^{-2\sigma}F(\zeta(\sigma)) = \zeta/F'(\zeta) - F(\zeta)/(F'(\zeta))^2$ and $\hat{g}$ is the determinant of $\hat{g}^{\mu\nu}$. In deriving Eqs. (2.22), we have used $e^\sigma = F'(\zeta)$. In addition, $\zeta(\sigma)$ is obtained by solving $e^\sigma = F'(\zeta)$ with respect to ζ as $\zeta = \zeta(\varphi)$. Defining φ as $\varphi \equiv \sqrt{3/2}\sigma/\kappa$, the action (2.22) is reduced to the following form of the canonical scalar field theory:

$$S_{\text{ST}} = \int d^4x \sqrt{-\hat{g}} \left[\frac{\hat{R}}{2\kappa^2} - \frac{1}{2}\hat{g}^{\mu\nu}\partial_\mu\varphi\partial_\nu\varphi - V(\varphi) + e^{-2\sqrt{2/3}\kappa\varphi}\mathcal{L}_{\text{matter}} \right]. \quad (2.23)$$

Taking the variation of the action (2.1) with respect to the metric $g_{\mu\nu}$, we find that the field equation of modified gravity is given by $F'(R)R_{\mu\nu} - (1/2)g_{\mu\nu}F(R) + g_{\mu\nu}\square F'(R) - \nabla_\mu\nabla_\nu F'(R) = \kappa^2 T_{\mu\nu}^{(\text{matter})}$. When there is no matter, using the $(\mu, \nu) = (0, 0)$ component and the trace part of the $(\mu, \nu) = (i, j)$ component of the above gravitational field equation, in the flat FRW background, we obtain

$$2\dot{H}F'(R) + 6\left(-4H^2\dot{H} + 4\dot{H}^2 + 3H\ddot{H} + \ddot{H}\right)F''(R) + 36\left(4H\dot{H} + \ddot{H}\right)^2 F'''(R) = 0. \quad (2.24)$$

We now investigate the case in which $F(R)$ is given by $F(R) = c_1 M^2 (R/M^2)^{-n}$, where c_1 is a dimensionless constant and M denotes a mass scale. The form of $F(R)$ in Eq. (2.19) corresponds to the above expression with $n = -7/2$. In this case, the scale factor $a(t)$ and the scale curvature R are given by [13] $a(t) = \bar{a}(t_s - t)^{(n+1)(2n+1)/(n+2)}$ and $R = 6n(n+1)(2n+1)(4n+5)(n+2)^{-2}(t_s - t)^{-2}$, respectively. It follows from $d\hat{t} = \pm e^{\sigma/2}dt$ that the relation between the cosmic time in the Einstein frame $\hat{t}$ and one in the Jordan frame is given by $\hat{t} = \mp\sqrt{-nc_1}(n+2)^n [6n(n+1)(2n+1)(4n+5)]^{-(n+1)/2} M^{n+1}(t_s - t)^{n+2}$. If $n < -2$, the limit of $t \rightarrow t_s$ corresponds to that of $\hat{t} \rightarrow \mp\infty$. For the case of Eq. (2.19), $n = -7/2$. Thus, we see that the Big Rip singularity does not appear in finite time for the scalar field theory, although it emerges in the corresponding modified gravitational theory. The metric in the Einstein frame is expressed as $d\hat{s}^2 = e^\sigma ds^2 = -d\hat{t}^2 + \hat{a}(\hat{t})d\mathbf{x}^2$, where $\hat{a}(\hat{t})$ is the scale factor in the scalar field theory given by $\hat{a}(\hat{t}) = \hat{a}\hat{t}^{3[(n+1)/(n+2)]^2}$, where $\hat{a}$ is a constant. For $n = -7/2$, because when $t \rightarrow t_s$, $\hat{t} \rightarrow \mp\infty$, the scale factor in the scalar field theory $\hat{a}(\hat{t})$ diverges at infinite time. Consequently, the (finite-time) Big Rip singularity in $F(R)$ gravity is transformed into the infinite-time singularity in the scalar field theory. This shows the physical difference of late-time cosmological evolutions between two mathematically equivalent theories.

III. FINITE-TIME FUTURE SINGULARITIES IN $F(R)$ GRAVITY

In this section, we examine several models of $F(R)$ -gravity with accelerating cosmological solutions ending at the finite-time future singularities by using the reconstruction technique explained in the preceding section.

First, we consider the case of the Big Rip singularity [17], in which H behaves as $H = h_0/(t_s - t)$. Here, h_0 and t_s are positive constants and H diverges at $t = t_s$. In this case, if we neglect the contribution from the matter, the general solution of (2.9) is given by

$$P(\phi) = P_+(t_s - \phi)^{\alpha_+} + P_-(t_s - \phi)^{\alpha_-}, \quad \alpha_\pm \equiv \frac{-h_0 + 1 \pm \sqrt{h_0^2 - 10h_0 + 1}}{2}, \quad (3.1)$$

when $h_0 > 5 + 2\sqrt{6}$ or $h_0 < 5 - 2\sqrt{6}$ and

$$P(\phi) = (t_s - \phi)^{-(h_0+1)/2} \left(A_1 \cos \left((t_s - \phi) \ln \frac{-h_0^2 + 10h_0 - 1}{2} \right) + B_1 \sin \left((t_s - \phi) \ln \frac{-h_0^2 + 10h_0 - 1}{2} \right) \right), \quad (3.2)$$

when $5 + 2\sqrt{6} > h_0 > 5 - 2\sqrt{6}$. Here, P_+ , P_- , A_1 and B_1 are constants. Using Eqs. (2.3), (2.4) and (2.10), we find that when R is large, the form of $F(R)$ is given by $F(R) \propto R^{1-\alpha_-/2}$ for $h_0 > 5 + 2\sqrt{6}$ or $h_0 < 5 - 2\sqrt{6}$ case and $F(R) \propto R^{(h_0+1)/4} \times (\text{oscillating parts})$ for $5 + 2\sqrt{6} > h_0 > 5 - 2\sqrt{6}$ case.

Next, we investigate more general singularity $H \sim h_0(t_s - t)^{-\beta}$ [18], where h_0 and β are constants, and h_0 is assumed to be positive and $t < t_s$ because it should be for the expanding universe. Even for non-integer $\beta < 0$, some derivative of H and therefore the curvature becomes singular. We should also note that in this case the scale factor a behaves as $a \sim \exp \left\{ [h_0/(\beta-1)](t_s - t)^{-(\beta-1)} + \dots \right\}$, where $\dots$ expresses the regular terms. From this expression, we find that if β could not be any integer, the value of a , and therefore the value of the metric tensor, would become complex number and include the imaginary part when $t > t_s$, which is unphysical. This could tell that the universe could end at $t = t_s$ even if β could be negative or less than -1 . We assume $\beta \neq 1$ because

TABLE I: Finite-time future singularities. Type I includes the case of ρ and p being finite at t_s . In case of Type IV, higher derivatives of H diverges. Type IV also includes the case in which p (ρ) or both of them tend to some finite values while higher derivatives of H diverge. Here, t_s is the time when a singularity appears and a_s is the value of $a(t)$ at $t = t_s$.

Type	Limit	a	ρ	$ p $
Type I (“Big Rip”)	$t \rightarrow t_s$	$a \rightarrow \infty$	$\rho \rightarrow \infty$	$ p \rightarrow \infty$
Type II (“sudden”)	$t \rightarrow t_s$	$a \rightarrow a_s$	$\rho \rightarrow \rho_s$	$ p \rightarrow \infty$
Type III	$t \rightarrow t_s$	$a \rightarrow a_s$	$\rho \rightarrow \infty$	$ p \rightarrow \infty$
Type IV	$t \rightarrow t_s$	$a \rightarrow a_s$	$\rho \rightarrow 0$	$ p \rightarrow 0$

the case $\beta = 1$ corresponds to the Big Rip singularity, which has been investigated. Furthermore, because the case $\beta = 0$ corresponds to de Sitter space, which has no singularity, we take $\beta \neq 0$. When $\beta > 1$, the scalar curvature R behaves as $R \sim 12H^2 \sim 12h_0^2(t_s - t)^{-2\beta}$. On the other hand, when $\beta < 1$, the scalar curvature R behaves as $R \sim 6\dot{H} \sim 6h_0\beta(t_s - t)^{-\beta-1}$. We may obtain the asymptotic solution for P when $\phi \rightarrow t_s$: (i) For $\beta > 1$, $P(\phi) \sim \exp\left\{[h_0/2(\beta-1)](t_s - \phi)^{-\beta+1}\right\}(t_s - \phi)^{\beta/2}\left(A_2 \cos\left(\omega(t_s - \phi)^{-\beta+1}\right) + B_2 \sin\left(\omega(t_s - \phi)^{-\beta+1}\right)\right)$, where $\omega \equiv h_0/[2(\beta-1)]$, and A_2 and B_2 are constants. When $\phi \rightarrow t_s$, $P(\phi)$ tends to vanish. (ii) For $1 > \beta > 0$, $P(\phi) \sim B_3 \exp\left\{-[h_0/2(1-\beta)](t_s - \phi)^{1-\beta}\right\}(t_s - \phi)^{(\beta+1)/8}$, where B_3 is a constant. (iii) For $\beta < 0$, $P(\phi) \sim A_3 \exp\left\{-[h_0/2(1-\beta)](t_s - \phi)^{1-\beta}\right\}(t_s - \phi)^{-(\beta^2-6\beta+1)/8}$, where A_3 is a constant. Using Eqs. (2.3), (2.4) and (2.10), we find the behavior of $F(R)$ (at large R) as summarized in Table II.

In the above investigations, we found the behavior of the scalar curvature R from that of H . Conversely, we now consider the behavior of H from that of R . When R evolves as $R \sim 6\dot{H} \sim R_0(t_s - t)^{-\gamma}$, if $\gamma > 2$, which corresponds to $\beta = \gamma/2 > 1$, H behaves as $H \sim \sqrt{R_0/12}(t_s - t)^{-\gamma/2}$, if $2 > \gamma > 1$, which corresponds to $1 > \beta = \gamma - 1 > 0$, H is given by $H \sim \{R_0/[6(\gamma-1)]\}(t_s - t)^{-\gamma+1}$, and if $\gamma < 1$, which corresponds to $\beta = \gamma - 1 < 0$, we obtain $H \sim H_0 + \{R_0/[6(\gamma-1)]\}(t_s - t)^{-\gamma+1}$, where H_0 is an arbitrary constant and it does not affect the behavior of R . H_0 is chosen to vanish in $H \sim h_0(t_s - t)^{-\beta}$. If $\gamma > 2$, we find $a(t) \propto \exp\left[(2/\gamma-1)\sqrt{R_0/12}(t_s - t)^{-\gamma/2+1}\right]$, when $2 > \gamma > 1$, $a(t)$ behaves as $a(t) \propto \exp\left(\{R_0/[6\gamma(\gamma-1)]\}(t_s - t)^{-\gamma}\right)$, and if $\gamma < 1$, $a(t) \propto \exp\left(H_0 t + \{R_0/[6\gamma(\gamma-1)]\}(t_s - t)^{-\gamma}\right)$. In any case, there appears a sudden future singularity [19] at $t = t_s$.

Since the second term in $H \sim H_0 + \{R_0/[6(\gamma-1)]\}(t_s - t)^{-\gamma+1}$ is smaller than the first one, we may solve Eq. (2.9) asymptotically as $P \sim P_0 \left\{1 + [2h_0/(1-\beta)](t_s - \phi)^{1-\beta}\right\}$ with a constant P_0 , which gives $F(R) \sim F_0 R + F_1 R^{2\beta/(\beta+1)}$, where F_0 and F_1 are constants. When $0 > \beta > -1$, we find $2\beta/(\beta+1) < 0$. On the other hand, when $\beta < -1$, we obtain $2\beta/(\beta+1) > 2$. As we saw in $F(R) \propto R^{1-\alpha-1/2}$ above, for $\beta < -1$, H diverges when $t \rightarrow t_s$. Since we reconstruct $F(R)$ so that the behavior of H could be recovered, the $F(R)$ generates the Big Rip singularity when R is large. Thus, even if R is small, the $F(R)$ generates a singularity where higher derivatives of H diverge.

We assume that H behaves as $H \sim h_0(t_s - t)^{-\beta}$. For $\beta > 1$, when $t \rightarrow t_s$, $a \sim \exp\left[h_0(t_s - t)^{1-\beta}/(\beta-1)\right] \rightarrow \infty$ and $\rho_{\text{eff}}, |p_{\text{eff}}| \rightarrow \infty$. If $\beta = 1$, we find $a \sim (t_s - t)^{-h_0} \rightarrow \infty$ and $\rho_{\text{eff}}, |p_{\text{eff}}| \rightarrow \infty$. If $0 < \beta < 1$, a goes to a constant but $\rho, |p| \rightarrow \infty$. If $-1 < \beta < 0$, a and ρ vanish but $|p_{\text{eff}}| \rightarrow \infty$. When $\beta < 0$, instead of $H \sim h_0(t_s - t)^{-\beta}$, one may assume $H \sim H_0 + h_0(t_s - t)^{-\beta}$. Hence, if $-1 < \beta < 0$, ρ_{eff} has a finite value $3H_0^2/\kappa^2$ in the limit $t \rightarrow t_s$. If $\beta < -1$ but β is not any integer, a is finite and ρ_{eff} and p_{eff} vanish if $H_0 = 0$ or ρ_{eff} and p_{eff} are finite if $H_0 \neq 0$ but higher derivatives of H diverge. We should note that the leading behavior of the scalar curvature R does not depend on H_0 in $H \sim H_0 + h_0(t_s - t)^{-\beta}$, and that the second term in this expression is relevant to the leading behavior of R . We should note, however, that H_0 is relevant to the leading behavior of the effective energy density ρ_{eff} and the scale factor a .

In Ref. [6], the finite-time future singularities has been classified as shown in Table I. The Type I corresponds to $\beta > 1$ or $\beta = 1$ case, Type II to $-1 < \beta < 0$ case, Type III to $0 < \beta < 1$ case, and Type IV to $\beta < -1$ but β is not any integer number. Thus, we have constructed several examples of $F(R)$ gravity showing the above finite-time future singularities of any type. It also follows from the reconstruction method that there appears Type I singularity for $F(R) = R + \tilde{\alpha}R^n$ with $n > 2$ and Type III singularity for $F(R) = R - \tilde{\beta}R^{-n}$ with $n > 0$, where $\tilde{\alpha}$ and $\tilde{\beta}$ are constants. In fact, however, even if some specific model contains the finite-time future singularity, one can always reconstruct the model in the remote past in such a way that the finite-time future singularity could disappear. Positive powers of the curvature (polynomial structure) usually help to make the effective quintessence/phantom phase become transient

TABLE II: Summary of the behavior of $F(R)$ gravity in case of $H \sim h_0 (t_s - t)^{-\beta}$. Here, $c_1 = [h_0/2(\beta - 1)](12h_0)^{-(\beta-1)/(2\beta)}$ and $c_2 = [h_0/2(1 - \beta)](-6h_0\beta)^{(\beta-1)/(\beta+1)}$. We note that $-6h_0\beta R > 0$ when $h_0, R > 0$.

Type I (“Big Rip”)		Type II (“sudden”)	Type III	Type IV
β	$\beta > 1$	$-1 < \beta < 0$	$0 < \beta < 1$	$\beta < -1, \beta : \text{not integer}$
$F(R)$	$F(R) \propto e^{c_1 R^{\frac{\beta-1}{2\beta}}} R^{-\frac{1}{4}}$	$F(R) \propto e^{-c_2 R^{\frac{\beta-1}{\beta+1}}} R^{\frac{\beta^2+2\beta+9}{8(\beta+1)}}$	$F(R) \propto e^{-c_2 R^{\frac{\beta-1}{\beta+1}}} R^{\frac{7}{8}}$	$F(R) \propto e^{-c_2 R^{\frac{\beta-1}{\beta+1}}} R^{\frac{\beta^2+2\beta+9}{8(\beta+1)}}$

and to avoid the finite-time future singularities. The corresponding examples have been examined in Refs. [3, 18].

IV. CONCLUSION

In the present article, we have reviewed finite-time future singularities in modified gravity. We have reconstructed an explicit model of modified gravity realizing a crossing of the phantom divide. It has been shown that the Big Rip singularity appears in this modified gravitational theory, whereas that the (finite-time) Big Rip singularity in the modified gravity is transformed into the infinite-time singularity in the corresponding scalar field theory. In addition, we have examined several models of modified gravity which predict accelerating cosmologies ending at the finite-time future singularities of all four known types.

Acknowledgments

The author deeply appreciates the invitation of Professor Sergei D. Odintsov to submit this article to the special volume *Casimir effect and Cosmology* on the occasion of the 70th birthday of Professor I. Brevik published by Tomsk State Pedagogical University, 2008. He also thanks Professor Chao-Qiang Geng, Professor Shin’ichi Nojiri and Professor S. D. Odintsov for their collaboration in Refs. [5, 9] very much. In addition, he is grateful to Professor Misao Sasaki for very helpful discussion of related problems. This work is supported in part by National Tsing Hua University under Grant #: 97N2309F1.

-
- [1] S. Nojiri and S. D. Odintsov, eConf **C0602061** (2006) 06 [Int. J. Geom. Meth. Mod. Phys. **4** (2007) 115] [arXiv:hep-th/0601213].
 - [2] S. Nojiri and S. D. Odintsov, arXiv:0801.4843 [astro-ph]; arXiv:0807.0685 [hep-th]; T. P. Sotiriou and V. Faraoni, arXiv:0805.1726 [gr-qc]; F. S. N. Lobo, arXiv:0807.1640 [gr-qc]; S. Capozziello and M. Francaviglia, Gen. Rel. Grav. **40** (2008) 357.
 - [3] M. C. B. Abdalla, S. Nojiri and S. D. Odintsov, Class. Quant. Grav. **22** (2005) L35 [arXiv:hep-th/0409177].
 - [4] L. Amendola and S. Tsujikawa, Phys. Lett. B **660** (2008) 125.
 - [5] K. Bamba, C. Q. Geng, S. Nojiri and S. D. Odintsov, arXiv:0810.4296 [hep-th].
 - [6] S. Nojiri, S. D. Odintsov and S. Tsujikawa, Phys. Rev. D **71** (2005) 063004 [arXiv:hep-th/0501025].
 - [7] S. Nojiri and S. D. Odintsov, Phys. Rev. D **74** (2006) 086005 [arXiv:hep-th/0608008].
 - [8] S. Nojiri and S. D. Odintsov, J. Phys. Conf. Ser. **66** (2007) 012005 [arXiv:hep-th/0611071]; J. Phys. A **40** (2007) 6725 [arXiv:hep-th/0610164].
 - [9] K. Bamba, S. Nojiri and S. D. Odintsov, JCAP **0810** (2008) 045 [arXiv:0807.2575 [hep-th]].
 - [10] J. L. Cortes and J. Indurain, arXiv:0805.3481 [astro-ph].
 - [11] S. Nojiri and S. D. Odintsov, Phys. Rev. D **77** (2008) 026007 [arXiv:0710.1738 [hep-th]]; G. Cognola, E. Elizalde, S. Nojiri, S. D. Odintsov, L. Sebastiani and S. Zerbini, *ibid.* **77** (2008) 046009 [arXiv:0712.4017 [hep-th]]; G. Cognola, E. Elizalde, S. D. Odintsov, P. Tretyakov and S. Zerbini, arXiv:0810.4989 [gr-qc].
 - [12] S. Nojiri and S. D. Odintsov, Phys. Lett. B **657** (2007) 238 [arXiv:0707.1941 [hep-th]].
 - [13] F. Briscese, E. Elizalde, S. Nojiri and S. D. Odintsov, Phys. Lett. B **646** (2007) 105 [arXiv:hep-th/0612220].
 - [14] S. Capozziello, S. Nojiri and S. D. Odintsov, Phys. Lett. B **634** (2006) 93 [arXiv:hep-th/0512118].
 - [15] Y. Fujii and K. Maeda, *The Scalar-Tensor Theory of Gravitation* (Cambridge University Press, Cambridge, United Kingdom, 2003); K. i. Maeda, Phys. Rev. D **39** (1989) 3159.
 - [16] S. Nojiri and S. D. Odintsov, Phys. Rev. D **68** (2003) 123512 [arXiv:hep-th/0307288].
 - [17] B. McInnes, JHEP **0208** (2002) 029.
 - [18] S. Nojiri and S. D. Odintsov, Phys. Rev. D **78** (2008) 046006 [arXiv:0804.3519 [hep-th]].

- [19] J. D. Barrow, *Class. Quant. Grav.* **21** (2004) L79; S. Cotsakis and I. Klaoudatou, *J. Geom. Phys.* **55** (2005) 306; S. Nojiri and S. D. Odintsov, *Phys. Rev. D* **70** (2004) 103522; J. D. Barrow and C. G. Tsagas, *Class. Quant. Grav.* **22** (2005) 1563; M. P. Dabrowski, *Phys. Rev. D* **71** (2005) 103505; *Phys. Lett. B* **625** (2005) 184; L. Fernandez-Jambrina and R. Lazkoz, *Phys. Rev. D* **70** (2004) 121503; **74** (2006) 064030; P. Tretyakov, A. Toporensky, Y. Shtanov and V. Sahni, *Class. Quant. Grav.* **23** (2006) 3259; M. Bouhmadi-Lopez, P. F. Gonzalez-Diaz and P. Martin-Moruno, *Phys. Lett. B* **659** (2008) 1; C. Cattoen and M. Visser, *Class. Quant. Grav.* **22** (2005) 4913; I. Brevik and O. Gorbunova, *Eur. Phys. J. C* **56** (2008) 425; J. D. Barrow, *Phys. Lett. B* **235** (1990) 40.